

830 Abdigermas copyediting

by MIT Id

Zinc-Chitosan Nanofibers for Necrotic Wound Bed Preparation in High-Grade Diabetic Foot Ulcers: A Prospective Case Series

Khairul Bahri*
Universitas Padjadjaran,
INDONESIA

Titis Kurniawan
Universitas Padjadjaran,
INDONESIA

Hartiah Haroen
Universitas Padjadjaran,
INDONESIA

31 Kana Fajar
Universitas Padjadjaran
INDONESIA

Article Info

Article history:

Received: July 24, 2026
Revised: August 26, 2026
Accepted: August 26, 2026

Keywords:

Diabetic Foot Ulcers;
DMIST Score;
Necrotic Tissue;
Wound-Bed Preparation;
Zinc-Chitosan Nanofiber.

Abstract

Background: Severe diabetic foot ulcers (DFUs) with extensive necrotic tissue require effective wound-bed preparation to reduce bioburden and promote granulation. Zinc-chitosan nanofibers possess antimicrobial, immunomodulatory and moisture balance properties that may enhance wound healing. However, clinical evidence for their use in high-grade necrotic DFUs remains limited.

Aims: This case series evaluated the effectiveness of zinc-chitosan nanofibers as a primary dressing for high-grade diabetic foot ulcers with extensive necrosis.

Methods: This prospective observational case series included three purposefully selected adults with high-grade DFUs containing $\geq 80\%$ slough. Wounds were assessed at baseline and week 1, 2 and 4 using the DMIST scoring system. Tissue perfusion was evaluated using the ankle brachial pressure index (ABPI), while laboratory investigations included blood glucose, glycated hemoglobin, serum albumin and leukocyte count. Standardized wound care comprised hypochlorous acid cleansing, staged autolytic and mechanical debridement, moisture balance optimization, zinc-chitosan nanofibers primary dressing, foam and multilayer secondary dressings, and culture-guided antibiotic therapy.

Result: Baseline DMIST score ranged from 23 to 24 reflecting severe inflammation, purulent exudate and 80 to 100% slough. ABPI value of 0.8 to 0.9 conformed adequate perfusion for debridement. By week 2, DMIST score decreased to 21 to 23 with reduce necrosis and early granulation. By week 4, all wounds achieved a DMIST score of 10, with 80 to 95% granulation tissue, marked reductions in exudate and infection, resolution of inflammation and early epithelization.

Conclusion: Zinc-chitosan nanofibers showed promising clinical benefit as a primary dressing accelerating wound-bed preparation in severe necrotic DFUs. Large controlled studies are needed to confirm efficacy and establish clinical recommendations.

9 cite this article: Author. (202X). Title. *Jurnal Ilmiah Pengabdian Masyarakat Bidang Kesehatan (Abdigermas)*, X(X), XX-XX.

This article is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/) ©202x by author/s

INTRODUCTION

8
The global prevalence of diabetes continues to rise steadily. In 2019, approximately 463 million people (9.3% of the world population) were living with diabetes. This number is projected to reach 570 million (10.2%) by 2030 and 700 million (10.9%) by 2045 (Saeedi et al., 2019). Indonesia faces a substantial burden of diabetes and its complication. Approximately 15% of people with diabetes are expected to develop diabetic foot ulcers (DFUs) (Yunir et al., 2023). With an estimates 10.3 million people living with diabetes, Indonesia ranks among the countries with the highest diabetes prevalence globally (Ceriello & Colagiuri, 2025). DFUs contribute substantially to disability and lower-limb amputation, accounting for up to 80% of diabetes-related amputations (Andhika, 2020). DFUs often present at an advanced stage because many patients remain underdiagnosed or seek care only after ulcer progression. Infection is a major clinical concern, with studies reporting infection in up to 50-60% of DFUs and a high proportion of patients presenting

* Corresponding author:

Khairul Bahri, Universitas Padjadjaran, INDONESIA. ✉ khairul24002@mail.unpad.ac.id

with severe ulcers. Approximately 70% of cases have been reported as Wagner grade >3, while infection may occur in up to 100% of patients with active DFUs (Armstrong et al., 2023; Sitompul et al., 2015).

Delayed intervention leads to impaired healing, increased complications, and frequent need for repeated debridement. Another report shows that approximately 18.2% of patients require multiple debridement procedures due to inadequate initial wound care (Jain & Viswanath, 2015). The challenge becomes even greater in high-grade DFUs dominated by necrotic or slough tissue. Heavy and persistent bioburden perpetuates the inflammatory phase, inhibits new granulation formation, and significantly delays healing (Atkin et al., 2019; Perrucini et al., 2021). This necessitates efficient and measurable wound-bed preparation strategies (Dayya et al., 2022). Nanomaterial-based technologies, including zinc-chitosan nanofibers, offer promising benefits for managing extensive necrosis by reducing bioburden, stimulating tissue regeneration, and optimizing moisture balance (Davies et al., 2025; Samuel et al., 2024). Zinc-chitosan has long been recognized as a wound-care material capable of accelerating tissue repair (Singh et al., 2017; Yadav et al., 2023). However, structured clinical data on the application of zinc-chitosan nanofibers as a primary wound-bed preparation agent in severe DFU cases with extensive necrosis remain scarce in international literature.

Severe DFUs often follow an unpredictable and prolonged healing trajectory¹⁶ making structured assessment tools essential for detecting meaningful changes over time. The DMIST scoring system which evaluates depth, maceration, infection/inflammation, size, tissue type, wound edges, and tunnelling and provides a sensitive and standardized framework for monitoring healing progression and predicting potential treatment failure at an early stage (Oe et al., 2020). Other studies also emphasize that structured wound assessment enhances clinical decision-making, reduces delays in intervention, and supports tailored treatment strategies (Jais & Pratama, 2023). Thus, DMIST is more than an assessment tool; it is a clinically meaningful method for understanding the patient's healing trajectory in a systematic and actionable way.

Therefore, this case series aimed to describe the clinical outcome of zinc-chitosan nanofibers dressing application in managing bioburden and necrotic tissue and promoting wound-bed improvement in three patients with high-grade diabetic foot ulcers (DFUs). In contrast to previous research focusing primarily on zinc-chitosan biomaterial fabrication and laboratory-based evaluation, this study evaluates its real-world clinical application as a primary dressing in three high-grade diabetic foot ulcer cases. Using DMIST scoring, the study documents wound-bed changes, particularly reductions in bioburden and necrotic tissue, and provides a structured clinical protocol for applying this nanofiber-based dressing in severe DFUs.

METHOD

¹ This study employed a prospective observational case series design involving three patients with high-grade diabetic foot ulcers (DFUs) presenting with extensive necrotic or slough tissue covering at least 80% of the wound bed and accompanied by clinical sign of local or systemic infection. A case series design was considered appropriate because the study aimed to provide an in-depth clinical description of wound-bed preparation and healing progression following the application of zinc-chitosan nanofibers in patients with severe DFUs, a clinical presentation that is relatively uncommon and frequently encountered in advanced wound-care practice. Three cases were included using consecutive case sampling during the study period. All patients presenting to the study site who met the predefined eligibility criteria were considered for inclusion until three eligible cases were enrolled. The small number of cases reflects the limited availability of patients who simultaneously met the specific clinical criteria, including high-grade ulcer severity, extensive necrotic tissue, active infection, adequate peripheral perfusion, and completion of the four-week treatment protocol. Cases were therefore not selected based on treatment response. The three patients had comparable baseline wound severity and fulfilled the same eligibility criteria, allowing consistent longitudinal observation of treatment response while limiting clinical heterogeneity.

Study was conducted at WOCARE Center Bogor, West Java, Indonesia. Ethical approval was obtained from the WOCARE Center Research and Development Committee (Approval No. 019/EC-WOCARE/VI/2026). Written informed consent was obtained from all participants before

enrollment and all patient information and clinical photographs were anonymized to maintaining confidentiality and protect participant privacy. Eligible participants were adults aged 50-63 years presenting with diabetic foot ulcers containing at least 80% necrotic or slough tissue, clinical evidence of local infection, adequate peripheral perfusion, and willingness to participate in the study and publication process. Patients were excluded if they had severe lower-limb ischemia indicated by an ankle-brachial pressure index (ABPI) below 0.5, osteomyelitis requiring surgical intervention, or were receiving immunosuppressive therapy, as these conditions could substantially alter wound healing and confound interpretation of treatment outcomes.

A commercially zinc-chitosan nanofibers was used as the primary dressing following standardized wound-bed preparation. Metcovazin Regular, a topical formulation zinc oxide and chitosan was used as the zinc-chitosan component, while Fibercelle, a gelling fiber dressing composed of sodium carboxymethyl cellulose (sodium-CMC), served as the fiber-based carrier. After wound cleansing and assessment, Metcovazin Regular was applied evenly to the Fibercelle dressing to coat dressing surface. The Metcovazin-coated Fibercelle was then placed directly onto the wound bed as the primary dressing, allowing the zinc-chitosan formulation to remain in contact with the wound surface while the fiber dressing provided absorption and structural support. This combined dressing was subsequently covered with an appropriate secondary dressing and changed according the predefined dressing protocol. Both materials were applied in their original commercial form without modification and directly to the prepared wound bed according to the standardized wound-care protocol. The dressing was applied and changed according to wound condition and exudate level throughout the four-week observation period. All participants received a standardized four-week wound-care protocol based in the principles of modern wound care. Wound management was delivered within a multidisciplinary service involving certified wound care nurses, general practitioners, and other healthcare professionals experienced in diabetic foot management. Tissue management consisted of autolytic and mechanical debridement to remove non-viable tissue and optimize the wound bed. Infection and inflammation were controlled through wound irrigation using hypochlorous acid (HOCL) with topical antimicrobial therapy administered when clinically indicated. Zinc-chitosan nanofibers were applied directly to the wound bed as the primary dressing and changed twice weekly throughout the treatment period and covered with a foam dressing as the secondary dressing to maintain an optimal moist wound environment and reduce the risk of peri-wound maceration. All procedures followed standardized protocols for wound cleansing, wound-bed preparation, dressing selection, and nursing assessment implemented in the clinical practice setting.

Clinical evaluation was performed at baseline and weekly throughout the four-week observation period using the validated DMIST scoring system, which assesses seven dimensions of wound healing, including Depth, Maceration Infection/Inflammation, Size, Tissue Type, Wound Edge and Tunnelling. Weekly DMIST assessment enabled objective monitoring of wound healing progression over time. Peripheral perfusion was evaluated using the ankle-brachial pressure index (ABPI), while laboratory parameters including leucocyte count, glycated hemoglobin (HbA1c), and random blood glucose levels were recorded at baseline and during follow-up to assess systemic infection status and glycemic control. Standardized digital wound photographs were obtained at each visit using consistent imaging procedures to provide visual documentation of tissue changes throughout treatment. "The primary outcome measures were reduction of necrotic tissue, increase in granulation tissue formation, improvement in DMIST scores, and resolution of clinical signs of infection. Because this investigation employed a case series design, data analysis focused on descriptive longitudinal evaluation rather than inferential statistical testing. Clinical findings from each patient were analyzed individually using within-case analysis to describe temporal changes in wound characteristics, DMIST scores, perfusion status, laboratory findings, and photographic documentation over the four-week treatment period. Subsequently, cross-case analysis was performed to compare similarities and differences in healing trajectories among the three patients, identify recurring clinical response patterns, and explore potential factors associated with successful wound-bed preparation following zinc-chitosan nanofiber application. This standardized assessment strategy enabled objective documentation of wound healing while providing a comprehensive description of the biological responses to zinc-chitosan nanofibers in preparing the wound bed high-grade necrotic diabetic foot ulcers.

RESULTS AND DISCUSSION

Results

Case Presentation 1

A 63-years-old man presented to the wound care clinic with a non-healing wound on the left foot (Figure Case 1). The ulcer was located in the metatarsal region and developed after accidental contact with a hot object, followed by inadequate initial wound management. The patient had poorly controlled type 2 diabetes mellitus, with an HbA1c of 13% and random blood glucose of 365 mg/dL. Initial laboratory findings showed hypoalbuminemia (3.1 g/dL) and marked leukocytosis ($26.7 \times 10^3/\mu\text{L}$), indicating compromised nutritional status and active systemic inflammation. The patient had previously received routine wound cleansing and dressing at another healthcare facility and was prescribed oral cefadroxil 500 mg three times daily. However, the wound continued to show persistent exudate and local inflammatory signs. His work required prolonged standing and mobility, which increased the importance of wound healing for returning to daily activities and work.



Figure Case 1: S 63-year-old man with a diabetic foot ulcer on the left metatarsal foot. Observation showed 95% granulation of the wound bed later (1c). The infection signs are reduced, such as improvement in oedema and erythema, lower exudate levels, and decreased malodor.

At the initial assessment, the ulcer showed extensive tissue loss and was not assigned a stage. The wound presented with erythema extending to the ankle, abundant purulent exudate, malodor, and 100% slough covering the wound bed. The foot also showed wet necrotic, edema, and violaceous discoloration on the dorsum, suggesting impaired local tissue perfusion. Wound culture identified *Escherichia coli*, and antibiotic therapy was subsequently adjusted to oral cefixime 200 mg twice times daily for three days according to the sensitivity results. Nutritional support included oral albumin supplementation (2x3 tablets/day), vitamin D and vitamin C.

A structured wound-care approach was implemented, including mechanical and autolytic debridement, wound cleansing with hypochlorous acid (HOCl), application of zinc-chitosan nanofiber as the primary dressing, and foam dressing as the secondary. The healing trajectory was monitored using the DMIST scoring system. The DMIST score decreased from 24 at week 1 to 10 at week 4, accompanied by reductions in wound depth and size, resolution of inflammatory signs, and progressive wound-bed improvement (Figure Case 1). The wound bed changed from 100% slough to approximately 95% health granulation tissue. Malodor resolved, purulent exudate decreased substantially, and early epithelialization was observed at the wound margins.

Case Presentation 2

60-years-old man presented to the wound care clinic with a persistent ulcer in the metatarsal region of the right foot that had developed following an automobile accident (Figure Case 2). Before receiving appropriate wound care, the patient had used conventional approaches, including

massage and herbal treatments, which may have contributed to additional tissue trauma and delayed healing. The patient appeared generally weak and had poorly controlled diabetes mellitus. Initial laboratory findings showed Leukocytosis ($14.1 \times 10^3/\mu\text{L}$), serum albumin (3.5 g/dL), HbA1c (13%), and a random blood glucose level of 283 mg/dL, indicated active inflammation, suboptimal nutritional status, and poor glycemic control. Wound culture identified *Escherichia coli*, and antibiotic therapy was adjusted to oral cefixime 200 mg twice times daily for seven days according to the susceptibility results. Supportive therapy included oral albumin supplementation (2x3 tablet/day), vitamin D and vitamin C.



Figure Case 2: H 60-year-old man with a chronic wound on the right metatarsal foot. Significant improvement was observed, such as complete granulation (2c). The infection signs showed considerable enhancement, such as reduced oedema and erythema.

The patient occupation required prolonged standing and regular physical activity, which represented a potential challenge to activity restriction and offloading during wound treatment. Prior to referral, he had received routine wound care with daily dressing changes; however, the wound showed limited clinical improvement. A structure wound-care protocol was subsequently implemented. Non-viable tissue was managed through mechanical and ¹¹olytic debridement, followed by wound cleansing with HOCL. Zinc-chitosan nanofibers were applied as the primary dressing, with foam dressing as the secondary layer to maintain an appropriate moist wound environment and protect the developing granulation tissue.

Wound progression was monitor using the DMIST score. The score decreased from 2¹ in week 1 to 23 in week 2 and subsequently to 10 in week 4. ¹is improvement was accompanied by a reduction in wound size and necrotic tissue, resolution of inflammatory signs, and progressive granulation tissue ¹rmation. The wound bed changed from 100% slough to approximately 95% granulation tissue, with no apparent clinical signs of infection at week 4 (Figure Case 2). Overall, the patient demonstrated substantial wound-bed improvement during the four-week multimodal wound-care protocol.

Case Presentation 3

A 50-years-old man presented to the wound care clinic with a three-week-old ³ulcer on the posterior aspect of the left foot (Figure Case 3). The wound developed after a foot injury while gardening. Before presentation, the patient managed the wound independently using rivanol and 0.9% NaCl compresses without professional wound assessment. At initial examination, the ulcer was deep and extensive, with thick yellow slough, purulent exudate, and surrounding inflammatory signs, indicating a chronic wound complicated by local infection. The patient appeared generally weak. Laboratory findings showed leukocytosis ($11.8 \times 10^3/\mu\text{L}$), hypoalbuminemia (3.3 g/dL), an HbA1c of 10%, and a random blood glucose level of 206 mg/dL, indicating poor glycemic control, active inflammation sigs, and suboptimal nutritional status. Wound culture identified *Escherichia coli*, and antibiotic therapy was adjusted to oral cefixime 200 mg twice times daily for seven days

according to the susceptibility results. Nutritional support included oral albumin supplementation and Vitamin D and C.




Figure Case 3: D 50-year-old man with a chronic wound on the left metatarsal foot. Significant improvement was observed, such as complete granulation and tendon exposure (3c). The infection signs showed considerable enhancement, such as reduced oedema and erythema.

The patient's occupation involved daily agricultural activities in wet environments and prolonged mobility, which presented potential challenges for wound protection, offloading and infection prevention. The depth and extent of the ulcer also impaired walking and occupational activities. Peripheral perfusion was assessed using the ankle-brachial index (ABI) with a value of 0.90 indicating adequate peripheral perfusion. Wound management consisted of mechanical and autolytic debridement to remove non-viable tissue, wound cleansing with HOCL, and application of c-chitosan nanofiber as the primary dressing. Foam dressing was used as the secondary dressing to maintain an appropriate moist wound environment and protect the developing tissue. The wound was assessed weekly using DMIST scoring.

The total DMIST score was 24 at week 1, decreased to 23 at week 2, and reached 10 by week 4. The improvement was accompanied by reduction in wound size and necrotic tissue, resolution of infection and inflammation and progressive development of granulation tissue. The wound-bed changes from 100% slough at baseline to a clean granulation with approximately 80% and visible clean tendon at week 4 (Figure Case 3). These findings indicate substantial wound-bed improvement during the four-week multimodal wound-care protocol.

Table 1. DMIST Score and Wound Progression

Case	Week 1	Week 2	Week 4	Clinical Wound Progression
Case 1	24	23	10	Week 1: Severe ulcer with 100% slough, heavy purulent exudate, malodor, edema, and extensive inflammation.
				Week 2: Slough 30% with tendon exposure, 70% early formation of granulation tissue, persistent inflammation, increased maceration, and the appearance of small tunneling.
				Week 4: Granulation reached 95%, slough 5%, no signs of infection, minimal odor and exudate, and wound edges showing early epithelialization.
Case 2	24	23	10	Week 1: 100% Large ulcer with thick necrotic tissue, purulent discharge, peri-wound inflammation, and positive <i>E. coli</i> culture.
				Week 2: Slough 90% and 10 granulation tissue in wound bed, Reduction in wound size, mild tunneling persisted, and inflammation remained present.

Case	Week 1	Week 2	Week 4	Clinical Wound Progression
				Week 4: Wound appeared clean, granulation reached 95%, slough 5%, infection resolved, and a stable wound environment was achieved.
Case 3	23	21	10	Week 1: 100% extensive ulcer with thick yellow slough, purulent discharge, intense inflammation, and irregular wound edges. Week 2: Slough 95% with tendon expose, presence of small tunneling, and early granulation formation of 5%. Week 4: Inflammation resolved, maceration controlled, granulation reached 80%, slough 20% and exposed tendon appeared healthier and ready for the proliferative phase.

Overall, the three cases demonstrated a consistent improvement in wound status throughout the four-week follow-up as reflected by a progressive reduction in DMIST score. Score decreased from 23-24 at baseline to 21-23 in week 2 and reached 10 by week 4. This improvement indicates a reduction in wound severity, necrotic burden, inflammation, and other adverse wound characteristics, accompanied by progressive granulation and stabilization of the wound bed. Although minor tunneling was observed in some cases during the healing process, it did not interfere with the overall positive wound progression. By week 4, the wounds had reached a more stable condition that supported the transition toward the proliferative of healing.




Figure 4. DMIST Score Progression

Overall, the three cases demonstrated an almost identical pattern of clinical improvement, reflected by a 13 –14-point reduction in DMIST scores within four weeks. This trend confirms that the combined use of HOCl cleansing, targeted debridement, moisture management, infection control, and zinc-chitosan nanofiber primary dressing provides consistent therapeutic benefits in high-grade diabetic foot ulcers (DFUs). The DMIST trajectory graph illustrates a steep and sustained decline across all cases, reinforcing the effectiveness and reproducibility of the implemented wound-care protocol.

Discussion

The findings case series demonstrated that zinc-chitosan nanofibers, incorporated into a standardized multidisciplinary wound-care protocols, were associated with substantial improvement in wound-bed preparation among patients with severe diabetic foot ulcers characterized by extensive necrotic tissue. Across all three patients, zinc-chitosan nanofibers consistently accelerated autolytic debridement, reduce wound bioburden, and promoted early granulation tissue formation during the first four weeks of treatment. Necrotic tissue burden decreased markedly from 80% to 100% at baseline to minimal residual necrosis by week four, while healthy granulation tissue increased to approximately 80% to 95% of the wound bed. Concurrently, clinical signs of inflammation and infection progressively resolved, indicating successful transition from the inflammation phase toward the proliferative phase of wound healing. Rather than indicating that zinc-chitosan nanofibers alone were responsible for these outcomes, the present findings suggest that their integration into a comprehensive wound-bed preparation strategy may facilitate biological process essential for tissue regeneration in complex DFUs. This observational is clinically important because extensive necrotic tissue is one of the principal barriers to healing and is strongly associated with persistent infection, delayed epithelialization, prolonged hospitalization and increased amputation risk.

The favorable healing trajectory observed in this series is biologically plausible and supported by recent advances in biomaterial research. Chitosan exhibit broad-spectrum antimicrobial activity, modulates macrophage polarization toward an anti-inflammatory phenotype, regulates cytokine production, scavenges reactive oxygen species, and stimulates fibroblast proliferation. Collagen deposition, angiogenesis, and extracellular matrix remodeling (Afrose et al., 2025; Ji et al., 2022; Wang et al., 2025). Experiment studies have further shown that incorporating zinc into chitosan-based materials can enhances antimicrobial, anti-inflammatory, antioxidant, and cell-migration properties (Mutlu et al., 2022). Electrospun chitosan /PVA/zinc oxide nanofibers have also demonstrated antibacterial and antioxidant activity with improve wound healing in diabetic wound models (Ahmed et al., 2018). More recently, zinc-ion coordinated chitosan/polycaprolactone composite nanofibers demonstrated enhanced antimicrobial and immunomodulatory activity, supporting the potential of zinc-containing chitosan nanofibers as multifunctional wound-care materials (Zhou et al., 2023). These findings provide biological support for clinical observations in the present case series, although direct extrapolation from experimental biomaterial studies to clinical effectiveness should be made cautiously. Incorporating zinc into the nanofiber matrix may further enhance keratinocyte migration, collagen synthesis, epithelial regeneration, antioxidant defense, and innate immune responses while maintaining structural integrity of the dressing. In additional, the porous nanofiber architecture provides an optimally moist and oxygen-permeable microenvironment that support cell migration and tissue repair. These complementary biological mechanisms likely explain the rapid reduction in necrotic tissue and consistent development of healthy granulation tissue observed across all three cases. Similar therapeutic benefits have been report in regional studies evaluating chitosan-based dressing for chronic DFUs, where reduced infection indices, accelerated granulation tissue formation, and shorter healing times were consistently observed (Bahri et al., 2024; Gitarja et al., 2018; Samuel et al., 2024; Shah et al., 2025).

Objective assessment using the DMIST instrument further strengthened the interpretation of wound healing progression. DMIST was developed specifically for monitoring diabetic foot ulcers across seven wound characteristics, including wound depth, tissue composition, inflammation, infection control, wound size, wound edges, and tunnelling. Validity has been demonstrated though correlations with established wound assessment instruments and its ability to discriminate wound severity and predict non-healing at four-weeks (Oe et al., 2020). In the present case series, the progressive reduction in DMIST score from 24, 24, and 23 at baseline to 10 by week four provided a standardized representation of changes in wound characteristics. The use of a validated multidimensional assessment tool minimized subjective interpretation and enable standardized longitudinal evaluation of healing progression, thereby strengthening clinical decision-making (Tsuchiya, 2023; Yotsu et al., 2023). However, these improvements should not be interpreted as evidence of the isolated efficacy of zinc-chitosan nanofibers. This interpretation is consistent with the principles of contemporary wound-bed preparation. The 2024 Delphi consensus on foot ulcer

management emphasizes systematic assessment and management of wound tissue, infection and inflammation, moisture balance, and other local and systematic factors as interconnected components of wound-bed preparation (Stacey et al., 2024). Instead, wound healing likely resulted from synergistic interactions between advanced biomaterial therapy and comprehensive multidisciplinary wound management. Patients received sequential wound-bed preparation consisting of autolytic debridement during the initial week, followed by combined autolytic and conservative sharp wound debridement (CSWD) during the second week, together with culture-guided antimicrobial therapy, glycemic optimization, nutritional support, and ³³intenance of an optimal moist wound environment (Afiani et al., 2019; Frykberg & Banks, 2015; Lipsky et al., 2020; Nair et al., 2024). Consequently, ^{the} observed clinical outcomes should be viewed as the result of an integrated therapeutic strategy rather than a single intervention. This approach is consistent with the IWGDF framework, which positions appropriate wound-bed preparation, debridement, basic wound dressing, and multidisciplinary standard care as the foundation of DFUs management before adjunctive wound-healing interventions are considered (Chen et al., 2024).

An additional factor that may have contributed substantially to wound healing was the incorporation of hypochlorous acid (HOCL) into the wound cleansing protocols. HOCL possesses rapid ²⁰antimicrobial activity against Gram-positive bacteria, Gram-negative bacteria, fungi and mature biofilms while demonstrating excellent tissue compatibility and minimal cytotoxicity toward viable cells (Burian et al., 2022; Serena et al., 2023). Previous clinical evaluation of super-oxidized solution in chronic wound also supports its potential role as a wound-cleansing agent for controlling wound bioburden and supporting wound management (Bahri, 2024). Reducing wound bioburden before application of zinc-chitosan nanofibers likely optimized the wound microenvironment and facilitated tissue regeneration. Current international guidelines similarly recommend integrating effective wound cleansing and infection control into multimodal DFU management due to successful wound-bed preparation requires simultaneous ²¹control of microbial burden, maintenance of moisture balance, and preservation of viable tissue (Armstrong et al., 2017; Hinchliffe et al., 2020; Lipsky et al., 2020). Therefore, the favorable outcomes observed in this case series probably reflect the combined effects of advances dressing technology and evidence-based wound-bed preparation principles.

Overall, the present findings are consistent with previous studies reporting beneficial effect of chitosan-based and metal-containing biomaterials in chronic wound management. In a recent case series from WOCARE Indonesia, nano-colloidal silver-zinc cream was associated with ³progressive infection control and complete wound closure in three with DFUs, although the follow-up period was substantially longer than that used in present study (Mokti et al., 2025). Clinical evidence is also emerging. In the CHITOWOUND randomized placebo-controlled trial, chitosan-based gel was associated with a greater median reduction in wound surface area than placebo after 10 weeks of treatment (Slivnik et al., 2024). Nevertheless, evidence regarding chitosan dressing remains heterogeneous. Several clinical investigations have demonstrated only modest improvements or no significant differences compared with other advanced wound dressing, particularly among patients with uncontrolled hyperglycemia, critical limb ischemia, persistent biofilm infection, extensive osteomyelitis or poor adherence to multidisciplinary treatment. Difference in wound severity, biomaterial composition, treatment duration outcome measure, and concomitant therapies may explain these discrepancies across studies. Compared with these reports, patients included in the present case series demonstrated preserved peripheral perfusion, standardized multidisciplinary care, and complete adherence to the treatment protocols, factors that may have enhanced the biological effectiveness of zinc-chitosan nanofibers and contributed to the consistent healing trajectories observed.

Preserved peripheral circulation also appeared to support favorable healing responses. ABPI values ranging from 0.8 to 0.9 confirmed adequate lower-extremity perfusion and suggested that delay wound healing was primary attributable to inflammatory, infectious, and metabolic disturbances rather than critical ischemia (Kim et al., 2023; Morozova et al., 2018). Consequently, optimizing of infection control and wound-bed preparation was likely sufficient to restore progression toward tissue regeneration. The reproducibility of healing trajectories across the three patients further suggests that this standardized treatment protocol may be feasible for implementation within specialized community-based wound-care services in Indonesia where

multidisciplinary management is available. These findings also have implementation of community-based wound care services. A standardized wound-bed preparation protocol that combines wound assessment, debridement, infection control appropriate dressing selection, nutritional support and glycemic management can provide a structured approach for managing complex DFUs outside tertiary hospital setting when appropriate clinical expertise and referral pathways are available. In community-based services, such as approach may support earlier identification of delay healing, facilities timely wound-bed optimization, and improve continuity of care for patients who require repeated wound assessment. The used of DMIST as structured assessment tool may further support consistent monitoring and communication among wound care professionals. Although the present case series cannot establish the effectiveness or cost-effectiveness of this model, the observed clinical trajectories provide preliminary evidence if its practical feasibility and generate a basic for further evaluation in larger community-based population. This implementation perspective is particularly relevant to wound-care and community health settings due to it emphasizes the translations of advanced dressing technology into structured, multidisciplinary clinical practice rather than an evaluation the biomaterial solely under experimental condition.

Despite these encouraging findings, several limitations should be acknowledged. This investigation involved only three patients treated at a single wound-care center using a case series design without a comparison group. Consequently, causal relationships cannot be attributed exclusively to zinc-chitosan nanofibers due to debridement, antimicrobial therapy, glycemic optimization, nutritional support and other components of multidisciplinary care may also have substantially contributed to wound healing. Furthermore, the relative short follow-up period precluded assessment of complete to wound closure, ulcer recurrence, functional recovery, and long-term durability of treatment outcomes. Therefore, the present findings should be regarded as preliminary clinical evidence that generates hypotheses rather than definitive proof of treatment efficacy.

Implications

The prospective clinical value of zinc-chitosan nanofibers as a primary for wound-bed preparation in high-grade DFUs with extensive necrotic tissue is illustrated in this case series. The integrated use of zinc-chitosan nanofibers in comprehensive wound care protocols may enhance early wound healing outcome as evidenced by the consistent reduction in DMIST score, resolution of infection and inflammation observed in all three cases. The result also lends credence to the utilization of structured wound assessment in conjunction with the DMIST scoring system to objectively monitor treatment response and inform clinical decision making. The method may aid the reduction of prolonged inflammation, the reduction of repeated debridement, and the earlier progression to the proliferation phase of wound healing in clinical practice.

Research contribution

This study offers preliminary clinical evidence supporting the use of zinc-chitosan nanofibers as primary dressing in high-grade DFUs that are distinguished by extensive necrotic. The case series, in contrast to previous research that primarily reported laboratory or biomaterial development, illustrates real world clinical implementation through the use of a standardized wound care protocol and serial DMIST assessment in a four-week period. The study also emphasizes the integration of nanofiber therapy with infection control, moisture support, perfusion assessment and staged debridement, thereby providing practical evidence for the management of modern DFUs particularly in resource-limited clinical setting.

Limitations

Several limitations should be acknowledged. First, the findings have limited generalizability because the case series included only three consecutive patients from a single wound-care center. Although consecutive case inclusion reduced investigator-driven selection, selection bias cannot be completely excluded because only patients who met the specific eligibility criteria and were treated at the study site were included. Second, the absence of a control group precludes direct comparison with standard wound care or other advanced wound dressings and limits the ability to determine whether the observed improvement was attributable specifically to zinc-chitosan nanofibers. Third,

the four-week follow-up period was relatively short and therefore did not allow assessment of complete wound closure, ulcer recurrence, functional recovery, or long-term treatment outcomes. Finally, the effect of zinc-chitosan nanofibers could not be isolated from the concurrent intervention, including mechanical and autolytic debridement, HOCL wound cleansing, antimicrobial therapy, nutritional supplement, glycemic management, and multidisciplinary wound care. Therefore, the observed improvements should be interpreted as preliminary clinical observations rather than definitive evidence of treatment efficacy. Larger prospective studies with appropriate comparison groups and longer follow-up are needed to confirm these findings.

Suggestions

Zinc-chitosan nanofibers dressing should be assessed in future research through RCTs that are adequately powered and have a more diverse patient population. In order to evaluate the cost-effectiveness, limb salvage, recurrence rates, and complete wound closure, a longer follow-up is necessary. Additionally, comparative studies with other advanced wound dressing are required to ascertain their relative clinical efficacy. Additionally, further research to investigate the biological mechanisms of zinc-chitosan nanofibers such as their impact on inflammation, angiogenesis, collagen deposited, and biofilm control, wound offers more robust evidence in support of their routine clinical application in the management of diabetic foot ulcers.

CONCLUSION

Zinc-chitosan nanofibers show strong potential as an effective wound-bed preparation agent for diabetic foot ulcers with extensive necrotic burden. When integrated into a comprehensive wound-care protocol including infection control, moisture management, and structured debridement these nanofibers contribute to substantial clinical improvement within 2-4 weeks. Combined with objective evaluation using DMIST, the intervention consistently produced marked reductions in necrosis, resolution of inflammation, and robust granulation formation across all three cases. This case series supports zinc-chitosan nanofibers as a promising therapeutic option for improving outcomes in severe DFUs, while underscoring the need for larger controlled studies to validate these findings across diverse clinical settings.

ACKNOWLEDGMENT

The researchers would like to thank the Magister Programs of the Faculty of Nursing Universitas Padjadjaran Sumedang and would like thanks to WOCARE Center, Bogor for the permittance for this study. Lastly, huge appreciations to the respondents who involved willingly this study.

AUTHOR CONTRIBUTION STATEMENT

KB, TK, HH and KF did conception of this study. Data Collections are done by KH and KF with disputes between the case study that were selected settled by KB and KF. KB and KF drafted the case study, while TK and HH did a critical revision of the case study. Final approval of the version to the published was granted by all authors.

AI DISCLOSURE STATEMENT

The author used AI technology (Google Gemini) during the preparation of this work for (grammar, refine sentence structure and translate). After using the tool/service, the author thoroughly reviewed and edited the content as needed and takes full responsibility for the content of the publication.

CONFLICTS OF INTEREST

The authors declare that there are no conflicts of interest, whether financial, personal or professional, with any party that could influence the objective, execution and result of this research.

REFERENCES

- Afiani, N., Santoso, S., N, T. H., & Yahya, M. F. N. (2019). Efektifitas Debridemen Mekanik Pada Luka Bakar Derajat III Terhadap Kecepatan Penyembuhan Luka. *Jkep*, 4(2), 93–103. <https://doi.org/10.32668/jkep.v4i2.254>
- Afrose, N., Chakraborty, R., & Rajendran, K. (2025). International Journal of Biological Macromolecules coatings for antibacterial and antibiofilm wound dressings: A review. *International Journal of Biological Macromolecules*, 320(P1), 145873. <https://doi.org/10.1016/j.ijbiomac.2025.145873>
- Ahmed, R., Tariq, M., Ali, I., Asghar, R., Noorunnisa Khanam, P., Augustine, R., & Hasan, A. (2018). Novel electrospun chitosan/polyvinyl alcohol/zinc oxide nanofibrous mats with antibacterial and antioxidant properties for diabetic wound healing. *International Journal of Biological Macromolecules*, 120, 385–393. <https://doi.org/https://doi.org/10.1016/j.ijbiomac.2018.08.057>
- Andhika, C. (2020). Clinical Profile of Diabetic Foot Amputation in Hasan Sadikin Hospital, Bandung, Indonesia. *Journal of Endocrinology, Tropical Medicine, and Infectious Disease (JETROMI)*, 2(2), 57–62. <https://doi.org/10.32734/jetromi.v2i2.3734>
- Armstrong, D. G., Boulton, A. J. M., & Bus, S. A. (2017). Diabetic Foot Ulcers and Their Recurrence. *The New England Journal of Medicine*, 376(24), 2367–2375. <https://doi.org/10.1056/NEJMra1615439>
- Armstrong, D. G., Tan, T.-W., M., A. J., & Bus., B. A. (2023). Diabetic foot ulcers: A Review. *Advance for Nurse Practitioners*, 17(7), 62–75. <https://doi.org/10.1001/jama.2023.10578>
- Atkin, L., Bučko, Z., Cutting, K., Moffatt, C., Probst, A., Romanelli, marco, S. Schultz, G., & Tettelbach, W. (2019). Implementing TIMERS: the race against hard-to-heal wounds Inflammation / infection Social factors Edge Regeneration Moisture Tissue. *Journal of Wound Care*. <https://doi.org/https://doi.org/10.12968/jowc.2019.28.Sup3a.S1>
- Bahri, K. (2024). IDF23-0556 Comparative Evaluation of Cleansing Agent Super-Oxidized Solution and Polyhexamethylene Biguanide 0,25% on Chronic Wound. *Diabetes Research and Clinical Practice*, 209, 111537. <https://doi.org/https://doi.org/10.1016/j.diabres.2024.111537>
- Bahri, K., Fajar, K., Nurani, I. A., & Hidayat, R. (2024). Integrating Zinc Cream-Chitosan and Cadexomer Iodine for Treating Infections in Diabetic Foot Ulcers: A Case Report. *Java Nursing Journal*, 2(2), 116–123. <https://doi.org/https://doi.org/10.61716/jnj.v2i2.40>
- Burian, E. . , Sabah, L., kirketerp- moller, K., Gundersen, G., & S.agren, M. (2022). Effect of Stabilized Hypochlorous Acid on Re-epithelialization and Bacterial Bioburden in Acute Wounds: A Randomized Controlled Trial in Healthy Volunteers. *Acta Demato Venerologica*, 1–7. <https://doi.org/10.2340/actadv.v102.1624>
- Ceriello, A., & Colagiuri, S. (2025). IDF global clinical practice recommendations for managing type 2 diabetes – 2025. In *Diabetes Research and Clinical Practice* (Vol. 222). <https://doi.org/10.1016/j.diabres.2025.112152>
- Chen, P., Vilorio, N. C., Dhatariya, K., Jeffcoate, W., Lobmann, R., McIntosh, C., Piaggese, A., Steinberg, J., Vas, P., Viswanathan, V., Wu, S., & Game, F. (2024). Guidelines on interventions to enhance healing of foot ulcers in people with diabetes (IWGDF 2023 update). *Diabetes/Metabolism Research and Reviews*, 40(3), e3644. <https://doi.org/https://doi.org/10.1002/dmrr.3644>
- Davies, D., Moody, A., Shrestha, S., & Zafar, B. (2025). RSC Advances Zinc-integrated PLGA / chitosan nano fi ber mesh : a platform for wound healing applications. *RSC Advances*, 15, 38846–38864. <https://doi.org/10.1039/D5RA03639A>
- Dayya, D., O'Neill, O., Habib, N., Moore, J., Iyer, K., & Huedo-Medina, T. B. (2022). Debridement of diabetic foot ulcers: public health and clinical implications - a systematic review, meta-analysis, and meta-regression. *BMJ Surgery, Interventions, and Health Technologies*, 4(1). <https://doi.org/10.1136/bmjst-2021-000081>
- Frykberg, R. G., & Banks, J. (2015). Challenges in the Treatment of Chronic Wounds. *Advances in Wound Care*, 4(9), 560–582. <https://doi.org/10.1089/wound.2015.0635>
- Gitarja, W. S., Jamaluddin, A., Wibisono, A. H., Megawati, V. N., & Fajar, K. (2018). Wound care management in Indonesia: issues and challenges in diabetic foot ulceration. *Wounds Asia*, 1(2), 13–17. <https://woundsasia.com/journal-articles/wound-care-management-in-indonesia->

- issues-and-challenges-in-diabetic-foot-ulceration/
- Hinchliffe, R. J., Forsythe, R. O., Apelqvist, J., Boyko, E. J., Fitridge, R., Hong, J. P., Katsanos, K., Mills, J. L., Nikol, S., Reekers, J., Venermo, M., Zierler, R. E., & Schaper, N. C. (2020). Guidelines on diagnosis, prognosis, and management of peripheral artery disease in patients with foot ulcers and diabetes (IWGDF 2019 update). *Diabetes/Metabolism Research and Reviews*, 36(S1). <https://doi.org/10.1002/dmrr.3276>
- Jain, A. K. C., & Viswanath, S. (2015). Debridement In Diabetic Foot Complications – An Analysis Of Debridement Using Amit Jain's Grading System For Debridement. *International Journal of Medical Science and Clinical Inventions*, 2(02), 761–765. <https://doi.org/https://doi.org/10.18535/ijmsci>
- Jais, S., & Pratama, K. (2023). Heliyon A diabetic foot wound healing assessment tool : A scoping review. *Heliyon*, 9(5), e15736. <https://doi.org/10.1016/j.heliyon.2023.e15736>
- Ji, M., Li, J., Wang, Y., Li, F., Man, J., Li, J., Zhang, C., Peng, S., & Wang, S. (2022). Advances in chitosan-based wound dressings : Modifications , fabrications , applications and prospects. *Carbohydrate Polymers*, 297(August), 120058. <https://doi.org/10.1016/j.carbpol.2022.120058>
- Kim, J., Nomkhondorj, O., An, C. Y., Choi, Y. C., & Cho, J. (2023). *Management of diabetic foot ulcers : a narrative review*. 40(4), 335–342. <https://doi.org/doi:10.12701/jyms.2023.00682>
- Lipsky, B. A., Senneville, É., Abbas, Z. G., Aragón-Sánchez, J., Diggle, M., Embil, J. M., Kono, S., Lavery, L. A., Malone, M., van Asten, S. A., Urbančič-Rovan, V., & Peters, E. J. G. (2020). Guidelines on the diagnosis and treatment of foot infection in persons with diabetes (IWGDF 2019 update). *Diabetes/Metabolism Research and Reviews*, 36(S1), 1–24. <https://doi.org/10.1002/dmrr.3280>
- Mokti, S. A., Gitarja, W. S., & Bahri, K. (2025). Enhancing diabetic foot ulcer management in WOCARE Indonesia: Impact of nano-colloidal silver-zinc cream on infection control and healing. *Diabetic Foot Journal*, 28(2), 39–46. <https://diabetesonthenet.com/diabetic-foot-journal/enhancing-diabetic-foot-ulcer-management-in-wocare-indonesia-impact-of-nano-colloidal-silver-zinc-cream-on-infection-control-and-healing/>
- Morozova, V. V., Kozlova, Y. N., Ganichev, D. A., & Tikunova, N. V. (2018). *Bacteriophage Treatment of Infected Diabetic Foot Ulcers*. 1693, 151–158. <https://doi.org/10.1007/978-1-4939-7395-8>
- Mutlu, N., Liverani, L., Kurtuldu, F., Galusek, D., & Boccaccini, A. R. (2022). Zinc improves antibacterial, anti-inflammatory and cell motility activity of chitosan for wound healing applications. *International Journal of Biological Macromolecules*, 213, 845–857. <https://doi.org/https://doi.org/10.1016/j.ijbiomac.2022.05.199>
- Nair, H., Subramanian, B., Frescos, N., Singh, R., Srisawasdi, G., Vaidya, S., & Wong, F. (2024). Autolytic continuous debridement with a focus on biofilm management: Consensus document for the APAC region. In Wounds International (Ed.), *Wounds International*. Wounds International. www.woundsinternational.com
- Oe, M., Yotsu, R. R., Mhs, D. A., Sakai, R. N. Y., Tamaki, T., & Sugama, J. (2020). Validity of DMIST for monitoring healing of diabetic foot ulcers. *Wound Repair and Regeneration*, 28(4), 539–546. <https://doi.org/10.1111/wrr.12816>
- Perrucini, de O., Oliveira, P. D. de, Medeiros, R. F. de, Bertin, F. B. P., Pires-Oliveira, L. D. de A., Rederico, D. A., & Poli, R. C. (2021). Ultrasonic therapy modulates the expression of genes related to neovascularization and inflammation in fibroblasts. *Fisioterapia Em Movimento*, 34, 1–10. <https://doi.org/10.1590/fm.2021.34112>
- Saeedi, P., Petersohn, I., Salpea, P., Malanda, B., Karuranga, S., Unwin, N., Colagiuri, S., Guariguata, L., Motala, A. A., Ogurtsova, K., Shaw, J. E., Bright, D., & Williams, R. (2019). Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9(th) edition. *Diabetes Research and Clinical Practice*, 157, 107843. <https://doi.org/10.1016/j.diabres.2019.107843>
- Samuel, D., Rajkumar, R., & Vijayaragavan, V. (2024). *Chitosan-Based Biomaterial in Wound Healing : A Review*. 16(2). <https://doi.org/10.7759/cureus.55193>
- Serena, T. E., Harding, K., & Queen, D. (2023). Point-of-care fluorescence imaging to optimise wound bed preparation prior to cellular and / or tissue-based product (CTP) application. *Internasional Wound Jurnal*, October, 3441–3442. <https://doi.org/10.1111/iwj.14446>
- Shah, J., Patel, D., Ranavavare, D., Hudson, D., Tran, M., Schloss, R., Langrana, N., Berthiaume, F., &

- Kumar, S. (2025). Recent Advancements in Chitosan-Based Biomaterials for Wound Healing. *Journal of Functional Biomaterials*, 16(2), 45. <https://doi.org/https://doi.org/10.3390/jfb16020045>
- Singh, R., Shitiz, K., & Singh, A. (2017). *Chitin and chitosan : biopolymers for wound management OH*. 1–14. <https://doi.org/10.1111/iwj.12797>
- Sitompul, Y., Budiman, B., Soebardi, S., & Abdullah, M. (2015). Profil Pasien Kaki Diabetes yang Menjalani Reamputasi di Rumah Sakit Cipto Mangunkusumo Tahun 2008 -2012. *Jurnal Penyakit Dalam Indonesia*, 2(1), 9. <https://doi.org/10.7454/jpdi.v2i1.75>
- Slivnik, M., Navodnik Preložnik, M., Fir, M., Jazbar, J., Čebren Lipovec, N., Locatelli, I., Liette Lauzon, H., & Urbančič Rovar, V. (2024). A randomized, placebo-controlled study of chitosan gel for the treatment of chronic diabetic foot ulcers (the CHITOWOUND study). *BMJ Open Diabetes Research & Care*, 12(3), e004195. <https://doi.org/10.1136/bmjdr-2024-004195>
- Stacey, M. C., Yifter, H., Gregory, J. H., Whiteside, C., Kotru, B., Jahnke, D., Hill, M., Nair, H., Chan, B., Harley, C., Schultz, G., Murphy, C., Orellana, C., Landis, S., Dofitas, B., Sibbald, R. G., Heil, J., Smart, H., Goodman, L., ... Armstrong, D. G. (2024). Wound Bed Preparation 2024: Delphi Consensus on Foot Ulcer Management in Resource-Limited Settings. *Advances in Skin & Wound Care*, 37(4), 180–196. <https://doi.org/10.1097/ASW.000000000000120>
- Tsuchiya, S. (2023). *Relationship between items of DMIST and healing of diabetic foot ulcers*. March 2022, 345–350. <https://doi.org/10.1111/iwj.13880>
- Wang, Z., Zeng, X., Feng, W., Lu, Y., & Wei, P. (2025). Reframing chronic wound therapy : from growth factor delivery to regenerative immuno-engineering. *Frontiers in Immunology*, 16(1683591), 1–8. <https://doi.org/10.3389/fimmu.2025.1683591>
- Yadav, M., Kaushik, B., Rao, G. K., Srivastava, C. M., & Vaya, D. (2023). Advances and challenges in the use of chitosan and its derivatives in biomedical fields : A review. *Carbohydrate Polymer Technologies and Applications*, 5(May), 100323. <https://doi.org/10.1016/j.carpta.2023.100323>
- Yotsu, R. R., Oe, M., Sanada, H., & Tamaki, T. (2023). *From prototype to implementation: Development of the DMIST scoring system for monitoring diabetic foot ulcers*. 1–6. <https://doi.org/10.35772/ghmo.2023.01024>
- Yunir, E., Waspadji, S., Juli, T., Tarigan, E., Sarumpaet, A., & Christabel, E. V. (2023). *Pedoman IWGDF mengenai Pencegahan dan Tatalaksana Kaki Diabetes* (E. Yunir (ed.)). PIP INTERNA. <https://iwgdfguidelines.org/wp-content/uploads/2025/02/Indonesian-translation-IWGDF-2019.pdf>
- Zhou, F., Sun, S., Cui, C., Li, X., Wu, S., Ma, J., Chen, S., & Li, C. M. (2023). Zinc ions and ciprofloxacin-encapsulated chitosan/poly(ε-caprolactone) composite nanofibers promote wound healing via enhanced antibacterial and immunomodulatory. *International Journal of Biological Macromolecules*, 253, 127086. <https://doi.org/https://doi.org/10.1016/j.ijbiomac.2023.127086>

ORIGINALITY REPORT

12%

SIMILARITY INDEX

7%

INTERNET SOURCES

7%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	European Wound Management Association. "EWMA 2026 Conference Abstracts", Journal of Wound Management Official journal of the European Wound Management Association, 2026 Publication	1%
2	Matje Meriaty Huru, Kamilus Mamoh, Diyan Maria Kristin, Agustina Abuk Seran et al. "The Effect of the Pre-Screening Questionnaire for Child Development on Increasing Parental Compliance in Monitoring Child Development", MEDICA (International Medical Scientific Journal), 2026 Publication	1%
3	hmp-websites.s3.amazonaws.com Internet Source	1%
4	ejournal.gomit.id Internet Source	1%
5	ejournal.1001tutorial.com Internet Source	1%
6	ihj.ideajournal.id Internet Source	1%
7	Khawaja H. Haider. "Stem Cell-Free Therapy - Engineering Exosomes and Secretomes for Next-Generation Therapeutics", CRC Press, 2026 Publication	<1%
8	Submitted to London School of Economics and Political Science Student Paper	<1%
9	e-journal.gomit.id Internet Source	<1%
10	ecca.org Internet Source	<1%
11	woundsinternational.com Internet Source	<1%
12	pmc.ncbi.nlm.nih.gov Internet Source	<1%

13	Sofija Pejкова, Gordana Georgieva, Savetka Paljoskovska Jordanova, Stefania Azmanova Mladenovska et al. "Dellon Decompression of the Tarsal Tunnel: An Effective Approach to Improving Blood Flow, Promoting Ulcer Healing and Recovery of Sensibility in Diabetic Patients", Journal of Plastic, Reconstructive & Aesthetic Surgery, 2025 Publication	<1 %
14	"Pearls and Pitfalls in Skin Ulcer Management", Springer Science and Business Media LLC, 2023 Publication	<1 %
15	"Management and Strategies for Wound Healing", Springer Science and Business Media LLC, 2026 Publication	<1 %
16	Submitted to Empire State College Student Paper	<1 %
17	Submitted to Institute of Tropical Medicine Student Paper	<1 %
18	Submitted to Griffth University Student Paper	<1 %
19	Submitted to Online Education Services Student Paper	<1 %
20	acervodigital.ufpr.br Internet Source	<1 %
21	ejournal.unisayogya.ac.id Internet Source	<1 %
22	peerj.com Internet Source	<1 %
23	sajp.co.za Internet Source	<1 %
24	e-dmj.org Internet Source	<1 %
25	www.jssm.org Internet Source	<1 %
26	Putri Fauzia Sukmawati, Rizki Hidayat, Naziyah Naziyah. "Analisis Asuhan Keperawatan Luka Kaki Diabetik pada Tn.I dan Ny.A dengan Penggunaan Zinc Cream dan Chitosan sebagai Balutan Primer di Wocare Center Bogor", JURNAL KREATIVITAS PENGABDIAN KEPADA MASYARAKAT (PKM), 2022 Publication	<1 %

27	dergipark.org.tr Internet Source	<1 %
28	diabetesonthenet.com Internet Source	<1 %
29	journals.ai-mrc.com Internet Source	<1 %
30	mdpi-res.com Internet Source	<1 %
31	pdfs.semanticscholar.org Internet Source	<1 %
32	pubs.rsc.org Internet Source	<1 %
33	theses.gla.ac.uk Internet Source	<1 %
34	theses.lib.polyu.edu.hk Internet Source	<1 %
35	www.jks.fikes.unsoed.ac.id Internet Source	<1 %
36	www.mdpi.com Internet Source	<1 %
37	Jahangir Moini, Matthew Adams, Anthony LoGalbo. "Complications of Diabetes Mellitus - A Global Perspective", CRC Press, 2022 Publication	<1 %
38	Ovy Ganesan, Dennis P. Orgill. "An Overview of Recent Clinical Trials for Diabetic Foot Ulcer Therapies", Journal of Clinical Medicine, 2024 Publication	<1 %

Exclude quotes
Exclude bibliography

Off
On

Exclude matches

Off